

Monday to Friday 8am - 4pm

Phone: 9217 1777

FAX: 9217 1788

Email: MPaCCS@bethesda.asn.au

Internet (access referral forms):

www.bethesda.asn.au/mpaccs

Name of patient: _____	
Date of birth: _____	<input type="checkbox"/> Male <input type="checkbox"/> Female
Name of facility: _____	
Facility phone number: _____	
Facility locality: <input type="checkbox"/> North <input type="checkbox"/> South	

Priority of this referral is: <input type="checkbox"/> within 1-2 days	<input type="checkbox"/> within 5 days	<input type="checkbox"/> more than 5 days
Remember MPaCCS IS NOT AN EMERGENCY SERVICE		
Date of referral: _____	Date client will be discharged from hospital: _____	
Referring hospital/facility: _____	Client aware of referral?	<input type="checkbox"/> Yes <input type="checkbox"/> No
Ward/unit: _____	Family aware of referral?	<input type="checkbox"/> Yes <input type="checkbox"/> No
Referred by: _____	Client aware of diagnosis & prognosis?	<input type="checkbox"/> Yes <input type="checkbox"/> No
Position: _____	If No, why not? _____	
Other palliative care service involved? (please name): _____	GP aware of referral ?	<input type="checkbox"/> Yes <input type="checkbox"/> No
	GP name: _____	
	GP phone: _____	Fax: _____
	GP RETAINS CLINICAL RESPONSIBILITY FOR PATIENT	

Does the person have a progressive life-limiting condition (malignant or non-malignant)?

Primary Diagnosis: _____

Does the person have symptoms that require specialist assessment/management?

Details: _____

Does your facility need support or education to manage optimally?

Details: _____

Does the person/family have psychosocial or spiritual needs that require specialist assessment?

Details: _____

Does your facility need support with advance care planning for this resident?

Details: _____

Have you discussed the patient's treatment wishes?

(ie resident and family/carer understanding of palliative care, including NFR, antibiotics, transfusions, radiotherapy etc)

Details: _____

Is the resident in the terminal phase (hours/days until death)? Yes No

If YES please call 08 9217 1777 to discuss this referral with an MPaCCS nurse or social worker.

PLEASE INCLUDE: current medication chart, recent hospital discharge summary, advance care plan

Interpreter required? Yes No Language: _____

Who is the patient's health/medical decision maker? patient Public Advocate other:

- MPaCCS can assist facilities & GPs with specialist medical & psychosocial assessment, care planning & case review.
- Referrals are accepted from any medical and nursing staff at metropolitan hospitals, mental health & disability services, Department of Corrective Services & residential aged care facilities.
- The Supportive & Palliative Care Indicators Tool at www.spict.org.uk can help you identify residents who may need or benefit from a palliative care referral.

An MPaCCS nurse will call you promptly to triage your referral.

Residents who are having life-prolonging treatment are not excluded from referral to palliative care.

MPaCCS IS NOT AN EMERGENCY SERVICE – AFTER HOURS/WEEKEND CASES CALL 1300 55 86 55



Supportive and Palliative Care Indicators Tool (SPICT™)

The SPICT™ is used to help identify people whose health is deteriorating. Assess them for unmet supportive and palliative care needs. Plan care.

Look for any general indicators of poor or deteriorating health.

- Unplanned hospital admission(s).
- Performance status is poor or deteriorating, with limited reversibility. (eg. The person stays in bed or in a chair for more than half the day.)
- Depends on others for care due to increasing physical and/or mental health problems.
- The person's carer needs more help and support.
- The person has had significant weight loss over the last few months, or remains underweight.
- Persistent symptoms despite optimal treatment of underlying condition(s).
- The person (or family) asks for palliative care; chooses to reduce, stop or not have treatment; or wishes to focus on quality of life.

Look for clinical indicators of one or multiple life-limiting conditions.

Cancer

Functional ability deteriorating due to progressive cancer.

Too frail for cancer treatment or treatment is for symptom control.

Dementia/ frailty

Unable to dress, walk or eat without help.

Eating and drinking less; difficulty with swallowing.

Urinary and faecal incontinence.

Not able to communicate by speaking; little social interaction.

Frequent falls; fractured femur.

Recurrent febrile episodes or infections; aspiration pneumonia.

Neurological disease

Progressive deterioration in physical and/or cognitive function despite optimal therapy.

Speech problems with increasing difficulty communicating and/or progressive difficulty with swallowing.

Recurrent aspiration pneumonia; breathless or respiratory failure.

Persistent paralysis after stroke with significant loss of function and ongoing disability.

Heart/ vascular disease

Heart failure or extensive, untreatable coronary artery disease; with breathlessness or chest pain at rest or on minimal effort.

Severe, inoperable peripheral vascular disease.

Respiratory disease

Severe, chronic lung disease; with breathlessness at rest or on minimal effort between exacerbations.

Persistent hypoxia needing long term oxygen therapy.

Has needed ventilation for respiratory failure or ventilation is contraindicated.

Other conditions

Deteriorating and at risk of dying with other conditions or complications that are not reversible; any treatment available will have a poor outcome.

Kidney disease

Stage 4 or 5 chronic kidney disease (eGFR < 30ml/min) with deteriorating health.

Kidney failure complicating other life limiting conditions or treatments.

Stopping or not starting dialysis.

Liver disease

Cirrhosis with one or more complications in the past year:

- diuretic resistant ascites
- hepatic encephalopathy
- hepatorenal syndrome
- bacterial peritonitis
- recurrent variceal bleeds

Liver transplant is not possible.

Review current care and care planning.

- Review current treatment and medication to ensure the person receives optimal care; minimise polypharmacy.
- Consider referral for specialist assessment if symptoms or problems are complex and difficult to manage.
- Agree a current and future care plan with the person and their family. Support family carers.
- Plan ahead early if loss of decision-making capacity is likely.
- Record, communicate and coordinate the care plan.